

Integrating Landscape and Metapopulation Modeling Approaches: Viability of the Sharp-tailed Grouse in a Dynamic Landscape

H. REŞİT AKÇAKAYA^{1*}, VOLKER C. RADELOFF², DAVID J. MLADENOFF² AND HONG S. HE³

¹ Applied Biomathematics, 100 North Country Road, Setauket, NY 11733, U.S.A.

² Department of Forest Ecology and Management, University of Wisconsin-Madison, Madison, Wisconsin, U.S.A.

³ School of Natural Resources, University of Missouri-Columbia, Columbia, MO 65211 U.S.A

Abstract: *The scale of natural resource management is currently changing from local or stand scale to the landscape scale. The lack of management experience at this scale, and the limited feasibility of experiments have increased the use of scenario modeling to analyze the effects of different management actions on focal species. However, current modeling approaches are poorly suited for the analysis of viability in dynamic landscapes. Demographic (e.g., metapopulation) models of species living in these landscapes do not incorporate the variability in spatial patterns of early successional habitats, and landscape models have not been linked previously to population viability models. We link a landscape model to a metapopulation model and demonstrate the use of this model by analyzing the effect of forest management options on the viability of the Sharp-tailed Grouse (*Tympanuchus phasianellus*) in the Pine Barrens region of northwestern Wisconsin (USA). This approach allows viability analysis based on landscape dynamics brought about by processes such as succession, disturbances, and silviculture. The landscape component of the model (LANDIS) predicts forest landscape dynamics in the form of a time series of raster maps. These maps are combined into a time series of patch structures, which form the dynamic spatial structure of the metapopulation component (RAMAS). The results showed that the viability of this species was sensitive to landscape dynamics and demographic variables such as fecundity and mortality. Ignoring the landscape dynamics gives overly optimistic results and results based only on landscape dynamics (ignoring demography) lead to a different ranking of the management options than the ranking based on the more realistic model incorporating both landscape and demographic dynamics. Thus, models of species in dynamic landscapes must consider both habitat and population dynamics simultaneously.*

Introduction

Dynamics of species and the risks of extinction and decline they face are determined by their demographic characteristics, the distribution and quality of their habitat, and the changes in these factors in response to various natural processes and human-caused threats. Ecologists use two main approaches to model these dynamic processes: landscape models and metapopulation models.

Landscape models may predict changes in structure and composition of the vegetation cover or more general changes in classes of land cover (Baker 1989). Models that focus on vegetation change vary in the ecological processes they simulate (e.g., succession, disturbance), their spatial resolution (cell size), and the extent or total landscape area they may simulate (10s to 1,000,000s of ha). Models of landscape change that emphasize vegetation change largely evolved from the early forest simulation

* email resit@ramas.com

models of the 1970s (Botkin 1972; Ek & Monserud 1974) and the first generation of models of spatial landscape change of the 1980s (e.g., Sklar et al. 1985). Today, a variety of approaches to spatial simulation of vegetation change exist (Mladenoff & Baker 1999).

Metapopulation models predict the future of species existing in multiple populations, for example in landscapes fragmented by natural or anthropogenic processes. In these models, the focus is the persistence or viability of a focal species in a given landscape, often with risk-based methods collectively known as population viability analysis (Akçakaya & Sjögren-Gulve 2000). These include occupancy models (reviewed by Sjögren-Gulve & Hanski 2000), age- or stage-structured models (reviewed by Akçakaya 2000a), and individual-based models (reviewed by DeAngelis & Gross 1992; Lacy 2000).

There have been previous efforts to link vegetation succession or landscape models to wildlife-habitat models (e.g., Smith 1986; Davis & DeLain 1986; Hyman et al. 1991), to estimate, for example, future changes in habitats of threatened species (Pausas et al. 1997; Curnutt et al. 2000). Spatially explicit metapopulation models have incorporated elements of a static landscape into models that simulate the dynamics of a metapopulation that occupies the landscape (e.g., Fahrig & Merriam 1985; Lefkovitch & Fahrig 1985; Lamberson et al. 1992; Akçakaya et al. 1995). Other metapopulation models have also incorporated aspects of a dynamic landscape, for example, by modeling the carrying capacity (or other measure of the quality) of patches to change deterministically, stochastically, or both (e.g., as a deterministic function of time since a stochastic disturbance event) (Pulliam et al. 1992; Lindenmayer & Possingham 1995a, b, 1996; Akçakaya & Atwood 1997; Stelter et al. 1997; Akçakaya & Raphael 1998; Gaff et al. 2000; Johst et al. 2002; Keith, in press; Kindvall & Bergman, in press) or by considering types of patches based on occupancy and suitability and modeling the transitions between these types as stochastic events (Keymer et al. 2000; Amarasekare & Possingham 2001; Wahlberg et al. 2002).

However, none of these models included a dynamic spatial structure or a mechanistic model of landscape dynamics. Thus, by a “dynamic landscape,” we mean not only a landscape in which the quality (or suitability) of the patches changes, but also one in

which the spatial structure (location and number) of the patches may change. By “landscape model,” we mean not only a model that predicts such changes, but also one that makes these predictions by explicitly modeling vegetation or succession dynamics based on life-history attributes of species such as shade and fire tolerance, and distance and method of seed dispersal. RAMAS GIS (defined below) is the only generic metapopulation-modeling program that incorporates dynamic spatial structure, including appearing, disappearing, merging, and splitting patches (Akçakaya 2001, 2002), but it does not include an explicit landscape model that simulates vegetation dynamics. Thus, no previous model has combined landscape and metapopulation models with dynamic spatial structure to estimate extinction risks, although the need for, and possibility of, such a combination was recognized long ago (Burgman et al. 1993: 215; Holt et al. 1995). Here, we integrated the landscape model LANDIS with the metapopulation model RAMAS. Both LANDIS and RAMAS are spatially dynamic (i.e., they model temporal changes in the spatial structure of the landscape and the metapopulation, respectively). Both models are generic; thus, we believe the integrated model we used is applicable to many species that inhabit dynamic landscapes. We demonstrate the use of this integrated model by analyzing the viability of the Sharp-tailed Grouse (*Tympanuchus phasianellus*) in a dynamic landscape.

The LANDIS model (Mladenoff et al. 1996; Mladenoff & He 1999) simulates forest change by modeling tree species as ten-year age classes. It models succession based on interactions among species life-history characteristics, site conditions, disturbance regime and management. Life-history characteristics include longevity, age of sexual maturity, shade and fire tolerance, and seed dispersal distance. Site conditions are encapsulated by “land types,” which can be derived from climate, physiography, and soil texture and organic matter content. The LANDIS model incorporates natural processes (fire, windthrow, succession and seed dispersal), as well as forest harvesting and allows many different silvicultural treatments to be modeled (Mladenoff & He 1999, He & Mladenoff 1999a; Gustafson et al. 2000). LANDIS has been used to model forest landscape dynamics in north temperate forests of the U.S. Lake States (He & Mladenoff 1999b), the Missouri Ozarks (Shifley et al. 2000), chaparral in California (Franklin et al. 2001), and

Finnish boreal forest (Pennanen & Kuuluvainen 2002), and to investigate the effects of climate warming on forest ecosystems (He et al. 1999). Detailed discussion of model design and implementation are provided elsewhere (e.g., Mladenoff & He 1999; He & Mladenoff 1999b).

The RAMAS GIS model (Akçakaya 2002) simulates metapopulation dynamics with an age- or stage-structured model for each subpopulation. It uses landscape data to define the spatial structure of the metapopulation from the perspective of the habitat requirements of the modeled species (Akçakaya 2000b). Habitat dynamics are modeled by deterministic changes in model parameters (such as carrying capacity, fecundity, and survival), which are specified as time series of these parameters. Such time series are calculated from a time series of habitat maps that describe the changes in the landscape. The types of spatial changes may include changing habitat quality or patch size, as well as appearing, disappearing, merging, or splitting habitat patches (Akçakaya 2001). RAMAS GIS has been used to analyze the effect of habitat structure on a variety of species (Root 1998; Williams et al. 1999; Kindvall 2000; USFWS 2001; Brown et al. 2001; Broadfoot et al. 2001; Akçakaya et al. *in press*).

Methods

In our integration of LANDIS and RAMAS, we used the landscape model to create a time series of raster maps that describe the tree species composition and age distribution at each decade (Fig. 1). We then combined these maps into a time series of patch structures by using a habitat suitability function for the Sharp-tailed Grouse. This time series of patch structures forms the dynamic spatial structure of the metapopulation, in which the patches may be changing at 10-year time steps in the patterns summarized above. We modeled the local dynamics of the Sharp-tailed Grouse with matrix models at an annual time step, incorporating the cyclic oscillations of this species, and random environmental and demographic variation. Each step is detailed below.

Study species

Sharp-tailed Grouse habitat is dominated by grasses and shrubs (Connelly et al. 1998). Although the species occupies a large range in central and

northern North America, its numbers have declined greatly in many parts of its range due to conversion of its steppe-grassland habitat to agriculture and other forms of habitat loss (Connelly et al. 1998). In Wisconsin Sharp-tailed Grouse range has contracted since presettlement times because of habitat lost to agricultural lands and forests due to fire suppression (Gregg and Niemuth 2000) and forest planting. Currently, a large proportion of the species' habitat in Wisconsin is in the Pine Barrens region, with scattered habitat patches in other parts of the state. The habitat of Sharp-tailed Grouse is both fragmented (patchily distributed) and dynamic (the number, size, and distribution of patches change over time). Hence, it is well suited for demonstrating our integration of a landscape and a metapopulation model.

Study area

The northwest Pine Barrens region of Wisconsin (USA) is located on a 450,000-ha outwash plain with predominantly coarse, sandy soils (Murphy 1931; Radeloff et al. 1998, 1999). These soils are prone to drought and are conducive to fires. This area experienced significant fire disturbance before European settlement, when jack pine (*Pinus banksiana* Lamb.) forest and savanna dominated the landscape. Fires were common at that time, and after logging and early settlement in the late 1800s-early 1900s. Failed farming and some reforestation have occurred, so some of the original open barrens habitat has been maintained. However, the landscape pattern has been significantly altered in the last 150 years ago (Radeloff et al. 1999). Local management agencies are particularly interested in the effects of forest harvests on landscape pattern in our study area because of the decline of open habitat species (such as the Sharp-tailed Grouse) that persisted in fire generated openings at pre-settlement times.

Landscape prediction under alternative management scenarios

Past forest management has created landscape patterns that differ significantly from those typical under natural disturbance regimes (Mladenoff et al. 1993; Wallin et al. 1996). It has been suggested that forest management should aim to mimic natural disturbance pattern (Hunter 1993; Wallin et al. 1994); however, much is still unknown about the

effects of different forest management scenarios on landscape patterns. To help fill this knowledge gap, we simulated the change in forest landscape of the Pine Barrens under several scenarios of silvicultural management.

Using LANDIS, we simulated eight scenarios, using one of two alternative values for each of three silvicultural parameters: mean clear-cut size, minimum age of jack pine for harvest, and proportion of the landscape under red pine (*P. resinosa* Ait.) management (Table 1). The minimum age limit for clear cuts in jack pine (40 or 60 years) represents the entire range of rotation lengths commonly used in jack pine management in the U.S. Lake States (Benzie 1977). The range of clear-cut sizes (4 to 256 ha) captures the entire range of what is common in jack pine management in the U.S. Lake States. Standard deviation of the clear-cut size was set to one-quarter of the mean. The nutrient-limited soils of the Pine Barrens limit the number of tree species that can be successfully grown commercially to largely jack pine and red pine. Red pine management has increased recently in the Pine Barrens to limit future outbreaks of jack pine budworm (*Christoneura pinus pinus*) by increasing tree species diversity in the landscape (Radeloff et al. 2003). However, red pine management operates with rotation lengths that are about twice as long as those for jack pine thereby significantly decreasing the amount of open areas present at any given time. Half the scenarios did not include management for red pine, and all stands were eligible for harvests when they approached the minimum stand age. The other half of the scenarios included red pine management on 50% of the landscape. In these scenarios, jack pine harvests would only affect red pine < 50 years old. In areas where red pine was < 50 years old, these would be harvested together with jack pine creating a clearcut. However, in areas with red pine > 50 years old, these older cohorts would not be harvested, and a jack pine harvest event would result in a thinning of the red pine, removing all competitors. Once red pine was 100 years old it became eligible for red pine harvesting. The details of these landscape simulations are described by Radeloff et al. (2003).

Sharp-tailed Grouse habitat dynamics

The results of a LANDIS simulation are a time series of raster maps describing the species composition and age structure of tree species in the landscape.

We converted each of the maps into a time series of habitat maps for the Sharp-tailed Grouse by using information about the species' habitat requirements.

Habitat requirements of the Sharp-tailed Grouse

Habitats used by the Sharp-tailed Grouse are dominated by relatively dense herbaceous cover and shrubs, but there is considerable variation in the species composition (Connelly et al. 1998). We defined optimal habitat as areas with no trees older than 10 years and areas dominated by herbaceous vegetation and suboptimal habitat as areas dominated by 10- to 20-year old jack pine. We assigned a habitat suitability (HS) value of 1.0 for optimal and 0.5 for suboptimal habitat.

In the input data maps for the LANDIS simulations, there are open areas in the initial landscape that are unrelated to harvesting. These are wildlife management areas, oldfields, and sedge marshes. In the LANDIS simulations these quickly convert to forest, although many of them are actually maintained as open areas. To account for this, we kept the actively maintained wildlife areas as optimal habitat (HS=1.0) throughout the simulation, regardless of what the LANDIS simulations predicted for these areas. In essence, we assumed that the current wildlife management areas would continue to be managed for the next 100 years.

Spatial structure of the metapopulation

In our study, LANDIS predicted raster maps of habitat at 100-m resolution. We converted these to habitat suitability (HS) maps as described above and averaged them into 500-m cell size. The HS values ranged from 0 (no habitat) to 1 (optimal habitat), and there was one HS map for each time step (decade) of LANDIS simulation. Based on this specification of habitat, RAMAS GIS identified the patch structure (size and location of habitat patches) each decade. This method of patch identification is described in Akçakaya et al. (1995) and Akçakaya (2000b).

After the habitat patches are identified, the program calculates the carrying capacity and initial number of individuals in each patch, based on a density function. To estimate this density function, we used data given by Evrard et al. (2000) on the area of different habitat types in Crex Meadows in the Wisconsin Pine Barrens (Table 2). We assumed that sedge marsh contributes half as much as optimal

habitat (grass and brush prairie) to habitat quality, pine/oak/aspens forest contributes 20%, and food lots (where additional food is provided) contribute twice as much. Thus, we calculated a total "habitat area" (equivalent to an area of optimal habitat) by multiplying the area of sedge marsh by 0.5, the area of grass or brush by 1.0, the area of pine/oak/aspens forest by 0.2, and the area of food lots by 2.0 (Table 2). Based on these assumptions, the total "habitat area" in Crex Meadows was about 55 km². In this area, there was an average of 73.2 displaying males/year between 1980 and 1999 (Evrard et al. 2000; Table 2). Assuming displaying males are over 2 years old and they make up 9.7% of the population (based on the stable age structure of the stage matrix; see below), this corresponds to an average of 755 birds (all ages, male and female). This gives an average density of 13.7 birds/km² or 7.3 ha/bird in optimal habitat. This calculation is further supported by the fact that the estimated density is the same number as the average fall population observed on prime habitat on Drummond Island, Michigan over a seven-year period (Johnsgard 1983).

With the 500-m cell size (or 25-ha cell area), the density in optimal habitat is 3.45 birds/cell. We multiplied the total habitat suitability in each patch by this number. Total habitat suitability is the sum of the habitat suitability (HS) values of all the cells in the patch. Thus, the product of total HS in each patch and the density in optimal habitat (3.45 birds/cell) gives the maximum number of birds in that patch. We used the calculated numbers as the carrying capacities of the patches. Following Temple (1992), we assumed a ceiling-type density dependence model.

Metapopulation model based on landscape dynamics

With the process summarized above, we obtained a time series of maps of the patch structure in the landscape. During the 100-year simulation, the number and size of patches changed, depending on the processes (such as timber harvest) modeled by LANDIS. The changes occurred because patches of suitable habitat appeared or existing patches merged because of timber harvest, or they disappeared or split into multiple patches because of forest growth. The program identified these changes and the lineage of the patches (see Akçakaya 2001 for details of this method). The result was a metapopulation model of shifting patches in a dynamic landscape. Into this

dynamic spatial structure, we added demographic structure, as described in the next section.

Sharp-tailed Grouse demographic model

We combined the habitat-based spatial structure of the metapopulation with an age- and sex-structured stochastic model of the Sharp-tailed Grouse populations. The model included six stages; three for each sex: F0, F1 and F2+ are 0-year-old females, 1-year-old females, and 2-year-old and older females, respectively. The corresponding male stages are M0, M1 and M2+. The mating system is polygynous, with the ratio of breeding males to breeding females estimated as 1:10 (Robel 1972 cited in Temple 1992). Thus, in the model, the fecundities are multiplied by the minimum of the number of females and 10 times the number of males. Thus, we assumed each male could mate with up to 10 females.

Survival rates

Survival rates are similar between the sexes (Robel et al. 1972). The probability of a newly hatched female surviving to breed at 1 year of age is 0.26 (Sisson 1976 cited in Temple 1992). We assumed all surviving females breed at age 1 and that males may potentially breed starting at age 2. Average annual adult survival in an unharmed population is estimated as 0.53 (Schroeder 1994 cited in Connelly et al. 1998). During 1983-85, hunting mortality in Wisconsin was estimated as 30% (Gregg 1990). Since then, a permit process was initiated and bag limits were imposed. There are no recent studies on hunting mortality, but these measures are likely to have reduced the hunting mortality. We added a hunting mortality of 25% to our model as "harvest" of all age and sex classes. We also ran simulations varying the hunting mortality from 20% to 30% to analyze the sensitivity of viability to hunting pressure.

Reproduction

Average brood size ranges from 4.1 to 9.4, and offspring per breeding female is 6.1 (Robel et al. 1972, Connelly et al. 1998). However, these figures may reflect observations made substantially after hatching (and hence may reflect part of the first-year mortality). For our model, an estimate of hatchlings

per female is needed (because survival estimates are available from hatching to next breeding).

Average clutch size is 11.8 eggs, and the proportion of eggs that hatch ranges from 87% to 92%, with an average of about 90% (Connelly et al. 1998). There is only one brood per season. Thus, hatchlings per female per year can be estimated as 10.6. However, clutch size is lower (9.4 to 11.6) in reneating attempts (when the first nest is destroyed). The proportion of breeders reneating and the proportion of nonbreeders (whose all nesting attempts fail) are unknown. Thus, as a compromise, we assumed 8.36 hatchlings per female per year, the average of the two estimates (6.1 and 10.6). We assumed a sex ratio of 1:1 at birth.

We assumed post-breeding census: first age class (age 0) consists of hatchlings, which breed by the next census. Thus, age-0 fecundity = $0.26 * 8.36 = 2.17$ and age 1+ fecundity = $0.53 * 8.36 = 4.43$. These fecundities are divided equally between female and male offspring (e.g., average age 1+ fecundity = 2.22 daughters per female; Table 3). We varied fecundity plus and minus 5% to analyze the sensitivity of results to the uncertainty in this parameter.

Population cycles

Sharp-tailed Grouse populations in Wisconsin have a 10-year cycle (Evrard 2000). The generation length of the species, based on the above estimates of survival and fecundity, is about 2.1 years. Thus, these cycles are not based on the age structure or density dependence of the species but most likely on its relationships with other species. Although the causes of these cycles are not fully known, and data are not sufficient to model the trophic interactions that might cause these cycles in the Sharp-tailed Grouse, it is reasonable to assume that the proximate causes of these cyclic changes in the abundance and age structure of any particular species are changes in its fecundity and survival rates. Thus, the effect can be realistically simulated with oscillating vital rates. We added cyclicality to the fecundities and survival rates by multiplying each stage matrix elements with a “relative vital rate” coefficient that changed with a 10-year period (Fig. 2). The product of these 10 coefficients was 1.0, thus, they do not affect the long-term growth rate of the population.

Environmental variability

There are several other environmental factors, independent of the 10-year cycle discussed above, that cause fluctuations in the Sharp-tailed Grouse populations. Mortality increases during severe winters, and reproductive success may be determined by soil moisture because of its influence on cover and food (Connelly et al. 1998). Temple (1992) used coefficients of variation (CV) of 40% for fecundity, 30% for juvenile survival, and 20% for adult survival. However, his model did not include population cycles, which add a lot of temporal variability. We used 30%, 20%, and 5% for the CVs of fecundity, juvenile survival and adult survival, respectively, and sampled these vital rates from lognormal distributions.

Dispersal

We used the dispersal-distance function $M_{ij} = a \exp(-D_{ij}/b)$, where M_{ij} is the proportion of individuals in the source population j that move to the target population i , a , and b are the function parameters, and D_{ij} is the distance between the two populations. To estimate the parameters, we used data from Robel et al. (1972:95; weighted average for the two locations). According to these data, the proportion of dispersers (individuals moving more than 0.8 km) is 0.582. The average distance traveled by a dispersing juvenile female is 18.96 km. Based on the juvenile female dispersal distance, we calculated b as $-\ln(0.5) * 18.96 = 13.14$ and a as $0.582/b = 0.0443$. Based on the average dispersal distance of the other stages, we calculated their relative dispersal rates as 0.33, 0.66, 0.44, and 1.0, for adult males, adult females, juveniles males, and juvenile females, respectively. We used the relative dispersal rates to calculate the proportion of each age/sex class dispersing, based on the dispersal-distance function.

Simulations

We ran nine simulations: eight simulations with timber harvest scenarios that resulted in dynamic landscape as described above and one simulation (scenario 0) that was based on the initial landscape (year 0 in LANDIS simulations). Thus, under scenario 0, the habitat structure and distribution observed for year 0 did not change for the duration of the simulation. We ran each RAMAS simulation for 1000 replications. The duration of the simulations were set at 100 years. The results of the

simulations were summarized in terms of risk of decline to a set of thresholds (500, 1000, and 1500 birds) within 100 years.

Results

The Sharp-tailed Grouse habitat was distributed throughout the Pine Barrens region, with larger patches of suitable habitat in the middle and south sections of the region (Fig. 3). Analysis of the initial landscape (year 0 in LANDIS simulations) with respect to the habitat requirements of the Sharp-tailed Grouse resulted in a total carrying capacity of 16400 birds, distributed among 18 habitat patches.

Different timber harvest scenarios caused different patterns of change in the landscape, including the amount of available habitat, measured in terms of the total carrying capacity (K) of all habitat patches (Fig. 4 & Table 1). Scenario 0 (which assumed no habitat change) resulted in the largest K . The four scenarios (4, 8, 12, and 16) with a large clearcut size resulted in a larger average and final K than the four scenarios with a small clearcut size (Table 1).

The risk of decline of the Sharp-tailed Grouse to 500, 1000, and 1500 birds within the next 100 years was different under different scenarios (Table 4). Risks were highest with scenarios 8 and 16 and lowest with scenarios 0 and 4. Other measures of viability, expected minimum abundance (McCarthy & Thompson 2001) and total abundance at the end of the simulation, gave similar results (Table 4). When landscape dynamics were ignored (scenario 0), the risk of decline to 1000 birds within 100 years was < 0.0005 . For scenarios with landscape dynamics, this risk ranged from 0.005 to 0.916 (Table 4).

The results of all scenarios simulating landscape dynamics (i.e., all except for scenario 0) were sensitive to the changes in the value of average fecundity (Fig. 5). Risk of decline to 1000 birds within 100 years ranged from 0.05 to 1.0 when fecundity was 5% lower than the medium fecundity, and it ranged from 0.0 to 0.49 when fecundity was 5% higher than the medium fecundity. The results were also sensitive to the hunting mortality. Changing the hunting mortality from 20% to 30% increased the risk of decline to 1000 birds from 0-0.05 to 0.57-1.0 for scenarios with landscape dynamics (Fig. 6).

Discussion

One of the important issues in linking two spatially explicit models is the correspondence between spatial and temporal scales of the two models. In our integration of RAMAS and LANDIS, we kept both the temporal and the spatial scales flexible in order to use the most appropriate scale for each model. The rate of landscape changes was far slower than that of grouse population dynamics; hence the landscape model time step was longer (decadal) than metapopulation model time step (annual). The 100-m spatial resolution of the landscape model is realistic for simulating vegetation dynamics at the stand level but it is unnecessarily high for simulating the grouse dynamics at the population level. Even at the individual level, the 500-m spatial resolution (25-ha cell size) we used would be appropriate, considering that the home range of this species in Wisconsin is up to 259 ha (Connelly et al. 1988). However, our model is a patch- or population-based model and each patch consisted of dozens to hundreds of cells, providing sufficiently fine detail.

Our results indicated that the viability of the Sharp-tailed Grouse metapopulation in the Pine Barrens region depends on both landscape dynamics and demographic variables such as fecundity and mortality. The timber harvest scenario affected the amount of suitable habitat for the species, how the habitat changed over time, and the viability of the species. However, these effects were not always correlated. For example, although scenario 12 was one of the scenarios with the largest amount of habitat (total carrying capacity), it was also one of the three worst scenarios in terms of viability (Tables 4 & 5). Similarly, scenario 9 resulted in the smallest amount of habitat (lowest average and final K), but was one of the better scenarios in terms of viability. Such results suggest that ranking management options only in terms of the habitat they provide for threatened or declining species (ignoring the demography of the species) may be misleading.

On the other hand, our results also suggest that ignoring changes in the landscape may also be misleading. The simulations that assumed no change in the habitat (scenario 0) resulted in the highest viability. In other words, ignoring landscape dynamics overestimated viability and gave results that were too optimistic compared with the more realistic simulations that incorporated landscape dynamics.

The reason for these results may be complex interactions between habitat change (appearance, disappearance, splitting, and merging of habitat patches over time) and the population change (growth and dispersal potential). Johst et al. (2002) demonstrated that long-term metapopulation persistence depends on rates of patch destruction and regeneration and on rates of population growth and colonization and argued that these four variables may have different time scales, violating the assumption of time-scale separation in occupancy type of metapopulation models. This is further complicated in cases where the spatial structure of the patches may also be dynamic (due to merging and splitting patches).

Methods that do not incorporate dynamics in both habitat and demography may give an incomplete assessment of the management options. This is likely to be especially important in species, such as the Sharp-tailed Grouse, that live in fragmented landscapes and are dependent on temporary habitat patches. The viability of such species depends on the balance between the rate of appearance and spatial arrangement of patches on the one hand and the reproductive capacity of the species on the other. Consequently, the viability of the species depends sensitively both on the spatiotemporal pattern of landscape changes resulting from different silvicultural practices (Table 4) and on demographic variables such as fecundity (Fig. 5) and mortality (Fig. 6). Thus, neither habitat dynamics nor demographic properties alone can predict whether a certain forest management plan will result in a viable population of Sharp-tailed Grouse. The only way such an assessment can be made is by considering both habitat dynamics and population dynamics simultaneously.

Our model represents the first attempt at establishing a spatially explicit link between the landscape and the metapopulation approaches, allowing an analysis of population viability based on landscape dynamics brought about by natural and human mediated processes such as succession, disturbances, and silviculture. In addition, the integration of two generic, spatially explicit models allows application of this approach to other cases of species living in fragmented and dynamic habitats. The most important challenge to the practical use of this approach in conservation and management of species in dynamic landscapes comes from obtaining sufficiently precise estimates of model parameters.

Future developments focusing on statistical methods of data analysis and parameter estimation would greatly enhance the usefulness of this approach.

Acknowledgements

The development of the methods described in this paper was supported by the U.S. Department of Agriculture (grant 00-33610-9437). This simulation of different forest management scenarios was supported by a Research Joint Venture Agreement between the University of Wisconsin-Madison and the North-Central Research Station of the U.S. Forest Service and by the Integrated Science Services Bureau of the Wisconsin Department of Natural Resources.

Literature Cited

- Akçakaya, H.R. 2000a. Population viability analyses with demographically and spatially structured models. *Ecological Bulletins* **48**:23-38.
- Akçakaya, H.R. 2000b. Viability analyses with habitat-based metapopulation models. *Population Ecology* **42**:45-53.
- Akçakaya, H.R. 2001. Linking population-level risk assessment with landscape and habitat models. *Science of the Total Environment* **274**:283-291.
- Akçakaya, H.R. 2002. RAMAS GIS: Linking landscape data with population viability analysis (version 4.0t). Applied Biomathematics, Setauket, New York.
- Akçakaya, H.R., and J.L. Atwood. 1997. A habitat-based metapopulation model of the California Gnatcatcher. *Conservation Biology* **11**:422-434.
- Akçakaya, H.R., M.A. McCarthy, and J. Pearce. 1995. Linking landscape data with population viability analysis: management options for the helmeted honeyeater. *Biological Conservation* **73**:169-176.
- Akçakaya, H.R., and M.G. Raphael. 1998. Assessing human impact despite uncertainty: viability of the northern spotted owl metapopulation in the northwestern USA. *Biodiversity and Conservation* **7**:875-894.
- Akçakaya H.R., and P. Sjögren-Gulve. 2000. Population viability analysis in conservation planning: an overview. *Ecological Bulletins* **48**:9-21.
- Akçakaya, H. R., M.A. Burgman, O. Kindvall, C. Wood, P. Sjögren-Gulve, J. Hatfield, and M.A. McCarthy, editors. *Species Conservation and Management: Case Studies*. Oxford University Press [in press].
- Akçakaya, H. R., D. J. Mladenoff, and H. S. He. 2003. RAMAS Landscape: integrating metapopulation

- viability with LANDIS forest dynamics model. User Manual for version 1.0. Applied Biomathematics, Setauket, New York.
- Amarasekare, P., and Possingham, H.P. 2001. Patch dynamics and metapopulation theory: the case of successional species. *Journal of Theoretical Biology* **209**:333-344.
- Baker, W. L. 1989. A review of models of landscape change. *Landscape Ecology* **2**: 111-133.
- Benzie, J.W. 1977. Manager's handbook for jack pine in the north-central states. General technical report NC-32. US Forest Service, North Central Research Station, St. Paul, Minnesota.
- Botkin, D.B., J.F. Janak, and J.R. Wallis. 1972. Some ecological consequences of a computer model of forest growth. *Journal of Ecology* **60**:849-872.
- Broadfoot, J.D., R.C. Rosatte, and D.T. O'Leary. 2001. Raccoon and skunk population models for urban disease control planning in Ontario, Canada. *Ecological Applications* **11**:295-303.
- Brook, B.W., J. J. O'Grady, A. P. Chapman, M. A. Burgman, H. R. Akçakaya, R. Frankham. 2000. Predictive accuracy of population viability analysis in conservation biology. *Nature* **404**:385-387.
- Brown D.K., A.A. Echelle, D.L. Propst, J.E. Brooks, and W.L. Fisher. 2001. Catastrophic wildfire and number of populations as factors influencing risk of extinction for Gila trout (*Oncorhynchus gilae*). *Western North American Naturalist* **61**: 139-148.
- Burgman, M.A., S. Ferson and H.R. Akçakaya. 1993. Risk assessment in conservation biology. Chapman and Hall, London.
- Connelly, J. W., M. W. Gratson, and K. P. Reese. 1998. Sharp-tailed Grouse (*Tympanuchus phasianellus*) No. 148 in A. Poole and F. Gill, editors. *The birds of North America*, Philadelphia, Pennsylvania.
- Curnutt, J.L., J. Comiskey, M.P. Nott, and L.J. Gross. 2000. Landscape-based spatially explicit species index models for everglades restoration. *Ecological Applications* **10**: 1849-1860
- Davis, F.W., D.M. Stoms, J. E. Estes, J. Scean, and J.M. Scott. 1990. An information systems approach to the preservation of biological diversity. *International Journal of Geographical Information Systems* **4**:55-78.
- Davis, L.S., and L.I. DeLain. 1986. Linking wildlife-habitat analysis to forest planning with ECOSYM. Pages 361-369 in Verner, J., M.L. Morrison and C.J. Ralph (eds.) *Wildlife 2000: Modeling Habitat Relationships of Terrestrial Vertebrates*. University of Wisconsin Press, Madison.
- DeAngelis, D. L., L.J. Gross, editors. 1992. Individual-based models and approaches in ecology: populations, communities and ecosystems. Chapman and Hall, New York.
- Ek, A. R., and R. A. Monserud. 1974. FOREST: A computer model for the growth and reproduction of mixed species forest stands. Research report A2635. College of Agricultural and Life Sciences. University of Wisconsin-Madison, Madison, Wisconsin.
- Evrard, J.O., J.E. Hoefler, and P.A. Kooiker. 2000. The history of sharp-tailed grouse in the Crex Meadows Wildlife Area. *The Passenger Pigeon* **62**:175-183.
- Fahrig, L. and G. Merriam. 1985. Habitat patch connectivity and population survival. *Ecology* **66**:1762-68.
- Franklin, F., Syphard, A.D., Mladenoff, D.J., He, H.S., Simons, D.K., Martin, R.P., Deutschman, D., and J. F. O'Leary. 2001. Simulating the effects of different fire regimes on plant functional groups in Southern California. *Ecological Modelling* **142**: 261-283..
- Gaff, H., D. L. DeAngelis L. J. Gross, R. Salinas, and M. Shorrosh. 2000. A dynamic landscape model for fish in the Everglades and its application to restoration. *Ecological Modelling* **127**: 33-52.
- Gregg, L., and N.D. Niemuth. 2000. The history, status, and future of the sharp-tailed grouse in Wisconsin. *The Passenger Pigeon* **62**:159-174.
- Gregg, L.E. 1990. Harvest rates of sharp-tailed grouse on managed areas in Wisconsin. Wisconsin Department of Natural Resources. Madison, Wisconsin. Research report 152. 13 pp.
- Gustafson, E.J., Shifley, S.R., Mladenoff, D.J., Nimerfro, K.K., and He, H.S. 2000. Spatial simulation of forest succession and timber harvesting using LANDIS. *Canadian Journal of Forest Research* **30**: 32-43.
- He, H.S., and Mladenoff, D.J. 1999a. The effects of seed dispersal on the simulation of long-term forest landscape change. *Ecosystems* **2**:308-319.
- He, H.S., and Mladenoff, D.J. 1999b. Spatially explicit and stochastic simulation of forest landscape fire disturbance and succession. *Ecology* **80**:81-99.
- He, H.S., Mladenoff, D.J., and Crow, T.R. 1999. Linking an ecosystem model and a landscape model to study forest species response to climate warming. *Ecological Modelling* **114**:213-233.
- Holt, R.D., S.W. Pacala, T.W. Smith, and J. Liu. 1995. Linking contemporary vegetation models with spatially explicit animal population models. *Ecological Applications* **5**:20-27
- Hunter, M.L. 1993. Natural fire regimes as spatial models for managing boreal forests. *Biological Conservation* **65**:115-120.
- Hyman, J., J. McAninch, and D.L. DeAngelis. 1991. An individual-based simulation model of herbivory in a heterogeneous landscape. Pages 443-478 in M.G. Turner and R.H. Gardner, editors. *Quantitative methods in landscape ecology*. Springer-Verlag, New York.

- Johnsgard, P.A. 1983. The grouse of the world. University of Nebraska Press, Lincoln, Nebraska.
- Johnson, M. P. 2000. The influence of patch demographics on metapopulations, with particular reference to successional landscapes. *Oikos* **88**:67-74
- Johst, K., R. Brandl, and S. Eber. 2002. Metapopulation persistence in dynamic landscapes: the role of dispersal distance. *Oikos* **98**:263-270.
- Keith, D. Population viability analysis of the endangered Australian heath shrub, *Epacris barbata*, subject to recurring fires and disease epidemics. In Akçakaya, H.R., M.A. Burgman, O. Kindvall, C. Wood, P. Sjögren-Gulve, J. Hatfield, and M.A. McCarthy, editors. *Species Conservation and Management: Case Studies*. Oxford University Press [in press].
- Keymer J. E., P. A. Marquet, J. X. Velasco-Hernandez, and S. A. Levin. 2000. Extinction thresholds and metapopulation persistence in dynamic landscapes. *The American Naturalist* **156**:478-494
- Kindvall, O. 2000. Comparative precision of three spatially realistic simulation models of metapopulation dynamics. *Ecological Bulletins* **48**:101-110.
- Kindvall, O., and K-O. Bergman. Woodland brown butterfly *Lopinga achine* in Sweden. In Akçakaya, H.R., M.A. Burgman, O. Kindvall, C. Wood, P. Sjögren-Gulve, J. Hatfield, and M.A. McCarthy, editors. *Species Conservation and Management: Case Studies*. Oxford University Press [in press].
- Lacy, R. C. 2000a. Considering threats to the viability of small populations with individual-based models. *Ecological Bulletins* **48**:39-51.
- Lamberson, R., R. McKelvey, B.R. Noon, and C. Voss. 1992. A dynamic analysis of northern spotted owl viability in a fragmented landscape. *Conservation Biology* **6**:505-512.
- Lefkovich, L.P., and L. Fahrig. 1985. Spatial characteristics of habitat patches and population survival. *Ecological Modelling* **30**:297-308.
- Lindenmayer, D. B., and H. P. Possingham. 1995a. Modelling the impacts of wildfire on viability of metapopulations of the endangered Australian species of arboreal marsupial, Leadbeater's possum, *Gymnobelideus leadbeateri*. *Forest Ecology and Management* **74**:197-222.
- Lindenmayer, D. B., and H. P. Possingham. 1995b. The conservation of arboreal marsupials in the montane ash forests of the central highlands of Victoria, South-Eastern Australia. VII. Modelling the persistence of Leadbeater's possum in response to modified timber harvesting practices. *Biological Conservation* **73**:239-257.
- Lindenmayer, D. B., and H. P. Possingham. 1996. Ranking conservation and timber management options for Leadbeater's Possum in southeastern Australia using population viability analysis. *Conservation Biology* **10**:235-251
- McCarthy, M. A., and C. Thompson. 2001. Expected minimum population size as a measure of threat. *Animal Conservation* **4**:351-355.
- Mladenoff, D. J., and H. S. He. 1999. Design and behavior of LANDIS, an object-oriented model of forest landscape disturbance and succession, In D. J. Mladenoff and W. L. Baker, editors. *Spatial modeling of forest landscape change: approaches and applications*. Cambridge University Press, Cambridge.
- Mladenoff, D. J., and W. L. Baker. 1999. *Spatial modeling of forest landscape change: approaches and applications*. Cambridge University Press, Cambridge.
- Mladenoff, D.J., G.E. Host, J. Boeder, and T.R. Crow. 1996. LANDIS: A spatial simulation model of forest landscape disturbance, management, and succession. Pages 175-179 in: *Second International Conference on Integrating GIS and Environmental Modeling*. NCGIA. Santa Barbara, CA, USA.
- Mladenoff, D.J., White, M.A., Pastor, J., and Crow, T.R. 1993. Comparing spatial pattern in unaltered old-growth and disturbed forest landscapes. *Ecological Applications* **3**:294-306.
- Murphy, R.E. 1931. Geography of northwestern pine barrens of Wisconsin. *Transactions of the Wisconsin Academy of Science, Arts and Letters* **26**:96-120.
- Pausas, J.G., M.P. Austin, I.R. Noble. 1997. A forest simulation model for predicting eucalypt dynamics and habitat quality for arboreal marsupials. *Ecological Applications* **7**: 921-933.
- Pennanen, J., and T. Kuuluvainen. 2002. A spatial simulation approach to natural forest landscape dynamics in boreal Fennoscandia. *Forest Ecology and Management* **164**: 157-175.
- Pulliam, H. P, J. B. Dunning, and J. Liu. 1992. Population dynamics in complex landscapes: a case study. *Ecological Applications* **2**:165-177.
- Radeloff, V. C., D. J. Mladenoff, E. J. Gustafson, R. M. Scheller, H. S. He, P. A. Zollner, and H. R. Akçakaya. 2003. Modeling forest harvest effects on landscape pattern in the Northwest Wisconsin Pine Barrens [in review].
- Radeloff, V. C., D. J. Mladenoff, H. S. He, and M. S. Boyce. 1999. Forest landscape change: The northwest Wisconsin Pine Barrens before European settlement and today. *Canadian Journal of Forest Research* **29**:1649-1659.
- Radeloff, V. C., D. J. Mladenoff, K. L. Manies, and M. S. Boyce. 1998. Analyzing forest landscape restoration potential: Pre-settlement and current distribution of oak in the northwest Wisconsin Pine Barrens. *Transactions of the Wisconsin Academy of Sciences, Arts and Letters* **86**:189-205.

- Robel, R. J., F. R. Henderson, and W. Jackson. 1972. Some Sharp-tailed Grouse population statistics from South Dakota. *Journal of Wildlife Management* **36**:87-98.
- Root, K.V. 1998. Evaluating the effects of habitat quality, connectivity, and catastrophes on a threatened species. *Ecological Applications* **8**:854-865.
- Shifley, S. R., F. R. Thompson, D. R. Larsen, and W. D. Dijak. 2000. Modeling forest landscape change in the Missouri Ozarks under alternative management practices. *Computers & Electronics in Agriculture* **27**:7-24.
- Sjögren-Gulve, P., and I. Hanski. 2000. Metapopulation viability analysis using occupancy models. *Ecological Bulletins* **48**:53-71.
- Sklar, F.H., R. Costanza, and J.W. Day, Jr. 1985. Dynamic spatial simulation modelling of coastal wetland habitat succession. *Ecological Modelling* **29**:261-281.
- Smith, T.M. 1986. Habitat-simulation models: integrating habitat-classification and forest- simulation models. Pages 389-393 in J. Verner, M.L. Morrison, and C.J. Ralph, editors. *Wildlife 2000: Modeling habitat relationships of terrestrial vertebrates*. University of Wisconsin Press.
- Stelter, C., M. Reich., V. Grimm, and C. Wissel. 1997. Modelling persistence in dynamic landscapes: lessons from a metapopulation of the grasshopper *Bryodemus tuberculata*. *Journal of Animal Ecology* **66**: 508-518.
- Temple, S.A. 1992. Population viability analysis of a sharp-tailed grouse metapopulation in Wisconsin. Pages 750-758 in D. R. McCullough and R. H. Barrett, editors. *Wildlife 2001: Populations*. Elsevier Publishers, London.
- U.S. Fish and Wildlife Service. 2001. Southwestern Willow Flycatcher Recovery Plan. Albuquerque, NM.
- Wahlberg, N., T. Klemetti, and I. Hanski. 2002. Dynamic populations in a dynamic landscape: the metapopulation structure of the marsh fritillary butterfly. *Ecography* **25**:224-232.
- Wallin, D. O., F. J. Swanson, and B. Marks. 1994. Landscape pattern response to changes in pattern generation rules: land-use legacies in forestry. *Ecological Applications* **4**:569-580.
- Wallin, D. O., F. J. Swanson, B. Marks, B., Cissel, J.H., and Kertis, J. 1996. Comparison of managed and pre-settlement landscape dynamics in forests of the Pacific Northwest, USA. *Forest Ecology and Management* **85**:291-309.
- Williams L.R., A.A. Echelle, C.S. Toepfer, M.G. Williams, and W.L. Fisher. 1999. Simulation modeling of population viability for the leopard darter (Percidae : *Percina pantherina*). *The Southwestern Naturalist* **44**:470-477.

Tables

Table 1. Silvicultural parameters of the timber harvest scenarios and average, final, and minimum values of the total carrying capacity (K) resulting from each scenario sorted in order of decreasing average carrying capacity.

Scenario*	Jack pine min age for harvest	Clear-cut size (ha)	Percent in red pine management	Average <i>K</i>	Final <i>K</i>	Minimum <i>K</i>
0	n/a	n/a	n/a	16400	16400	16400
4	40	259	0	14544	14607	8939
12	60	259	0	10436	8727	5141
8	40	259	50	8031	5050	5050
16	60	259	50	7475	4878	4792
5	40	4	50	5809	5386	4570
13	60	4	50	5794	5381	4565
1	40	4	0	5764	4839	4340
9	60	4	0	5743	4812	4334

* Scenario 0 assumes a static landscape with no harvest, forest growth, fire, or any other landscape change. For the other scenarios, we followed the numbering in Radeloff (2003).

Table 2. Sharp-tailed Grouse Habitat in the Crex Meadows

Land type	Area ^a (km ²)	Habitat (%) ^b	Habitat area (km ²)
Open water	15.99	0	0.0
Sedge marsh	38.85	50	19.4
Grass/brush	29.95	100	29.9
Pine/oak/aspens	22.50	20	4.5
Food lots	0.51	200	1.0
Total	107.79		54.9

^a Evrard et al (2000)

^b Contribution of each land type to optimal habitat; see text for details.

Table 3. Average stage matrix used in the demographic model. The model includes three stages for each sex: F0, F1 and F2+ are 0-year-old females, 1-year-old females, and 2-year-old and older females, respectively. The corresponding male stages are M0, M1 and M2+.

	F0	F1	F2+	M0	M1	M2+
F0	1.0868	2.2154	2.2154	0	0	0
F1	0.26	0	0	0	0	0
F2+	0	0.53	0.53	0	0	0
M0	1.0868	2.2154	2.2154	0	0	0
M1	0	0	0	0.26	0	0
M2+	0	0	0	0	0.53	0.53

Table 4. Viability of Sharp-tailed Grouse in the Pine Barrens under nine scenarios of landscape change, in terms of risk of decline, expected minimum abundance and percentiles of abundance in year 100. The scenarios are sorted in order of increasing risk of decline.

Scenario	Probability of decline to threshold T within 100 years			Expected minimum total abundance	Percentiles of distribution of the total number of individuals in year 100				
	$T=500$	$T=1000$	$T=1500$		5th	25 th	50 th	75th	95th
0	0.000	0.000	0.000	5385	5409	6938	8200	9337	10793
4	0.000	0.005	0.064	2408	2033	3162	4178	5295	6919
13	0.000	0.017	0.573	1454	1787	2190	2500	2874	3314
5	0.000	0.020	0.559	1453	1789	2198	2522	2889	3399
9	0.000	0.082	0.818	1299	1221	1623	1929	2319	2774
1	0.000	0.099	0.833	1282	1143	1558	1896	2234	2754
12	0.013	0.256	0.768	1240	1412	1787	2056	2338	2924
8	0.150	0.906	1.000	720	388	618	779	990	1272
16	0.226	0.916	1.000	670	660	1136	1525	1889	2477




Figure 1. The integration of the landscape model LANDIS (Mladenoff et al. 1996) and the metapopulation model RAMAS (Akçakaya 2002) to model metapopulation viability in a dynamic forest landscape (Source: Akçakaya et al. 2003; reprinted by permission).




Figure 2. The time series of relative vital rates (multipliers for survival and fecundity) that were used to model population cycles in the Sharp-tailed Grouse model. There are 10 multipliers in each cycle of 10 years. The product of the 10 numbers is 1.00.




Figure 3. Distribution of sharp-tailed grouse habitat in the Wisconsin Pine Barrens region at year 0, 20, 40, and 80 (from top to bottom), under scenario 4 (Table 1).




Figure 4. Change in the total carrying capacity of all habitat patches (in terms of the number of grouse) with eight timber harvest scenarios.




Figure 5. Effect of Sharp-tailed Grouse fecundity on the species' risk of decline to 1000 birds within 100 years. Low and high fecundity values are 5% lower and higher than those described in the text, respectively.




Figure 6. Effect of hunting mortality on the risk of decline to 1000 birds within 100 years for scenarios 4, 5, 12, and 16 (Table 1). The results with the scenarios not shown here fell between the results for scenarios 4 and 16.